

INFUSION LOGBOOK

Timothy has severe hemophilia B and uses Rebinyn®.

A COMPREHENSIVE LOGBOOK TO KEEP YOUR BLEED AND INFUSION INFORMATION TOGETHER AND ORGANIZED

NAME: _____

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Indications and Usage

What is Rebinyn®, Coagulation Factor IX (Recombinant), GlycoPEGylated?

Rebinyn® is an injectable medicine used to replace clotting Factor IX that is missing in patients with hemophilia B. Rebinyn® is used to treat, prevent, or reduce the frequency (number) of bleeding episodes in people with hemophilia B. Your healthcare provider may give you Rebinyn® when you have surgery. Rebinyn® is not used for immune tolerance therapy.

Important Safety Information

What is the most important information I need to know about Rebinyn®?

- **Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center.** Carefully follow your healthcare provider's instructions regarding the dose and schedule for infusing Rebinyn®.

rebinyn®Coagulation Factor IX
(Recombinant), GlycoPEGylated

Tracking your bleeds and infusions is important. Keeping accurate records will help when you talk with your healthcare provider about your treatment. Together, you can have an informed conversation. Be sure to keep this logbook for future reference during your Rebinyn® journey.

2023

CIRCLE YOUR PROPHYLAXIS INFUSION DATES BELOW.

JANUARY						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

FEBRUARY						
S	M	T	W	T	F	S
		1	2	3	4	
5	6	7	8	9	10	11
12	13	14	15	16	27	18
19	20	21	22	23	24	25
26	27	28				

MARCH						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

APRIL						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

MAY						
S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

JUNE						
S	M	T	W	T	F	S
		1	2	3		
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	

JULY						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

AUGUST						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

SEPTEMBER						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

OCTOBER						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

NOVEMBER						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

DECEMBER						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

2024

CIRCLE YOUR PROPHYLAXIS INFUSION DATES BELOW.

JANUARY						
S	M	T	W	T	F	S
1	2	3	4	5	6	
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

FEBRUARY						
S	M	T	W	T	F	S
			1	2	3	
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29		

MARCH						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

APRIL						
S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

MAY						
S	M	T	W	T	F	S
		1	2	3	4	
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

JUNE						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

JULY						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

AUGUST						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

SEPTEMBER						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

OCTOBER						
S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

NOVEMBER						
S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

DECEMBER						
S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

Timothy has severe hemophilia B and uses Rebinyn®.

Please see additional Important Safety Information on page 11.
Please see Prescribing Information [here](#).



Bleed and treatment infusion records

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

This logbook may last for a longer period of time,
based upon your number of bleeds.



Abdiel has hemophilia B and uses Rebinyn®.



Bleed and treatment infusion records

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

HELPFUL TERM:

FACTOR LEVELS

In people with hemophilia, a clotting factor is missing or the level of factor is low. Hemophilia can be mild, moderate, or severe, depending on the level of clotting factor in a person's blood.

Please see additional Important Safety Information on page 11. Please see Prescribing Information [here](#).

This logbook may last for a longer period of time, based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

HELPFUL TERM:

MILD HEMOPHILIA

Hemophilia defined by Factor 9 levels
between 5% and 40%.

Please see Important Safety Information on page 11.
Please see Prescribing Information [here](#).

This logbook may last for a longer period of time,
based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated



Emili with her son Xander, who has hemophilia B and uses Rebinyn®.



Timothy has severe hemophilia B and uses Rebinyn®.



Bleed and treatment infusion records

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

HELPFUL TERM:

PROPHYLAXIS

The regular administration of factor to prevent bleeds.

Please see additional Important Safety Information on page 11. Please see Prescribing Information [here](#).

This logbook may last for a longer period of time, based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

HELPFUL TERM:

MODERATE HEMOPHILIA

Hemophilia defined by Factor 9
levels between 1% and 5%.

Please see Important Safety Information on page 11.
Please see Prescribing Information [here](#).

This logbook may last for a longer period of time,
based upon your number of bleeds.

rebinyn[®]

Coagulation Factor IX
(Recombinant), GlycoPEGylated



Abdiel has hemophilia B
and uses Rebinyn[®].



Timothy has severe hemophilia B and uses Rebinyn®.



Bleed and treatment infusion records

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

HELPFUL TERM:

HALF-LIFE

The amount of time it takes for the factor level to drop by half after an infusion.

Please see additional Important Safety Information on page 11. Please see Prescribing Information [here](#).

This logbook may last for a longer period of time, based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

HELPFUL TERM:

SEVERE HEMOPHILIA

Hemophilia defined by Factor 9 levels
less than 1%.

Please see Important Safety Information on page 11.
Please see Prescribing Information [here](#).

This logbook may last for a longer period of time,
based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated



Emili with her son Xander, who has hemophilia B and uses Rebinyn®.



Abdiel has hemophilia B and uses Rebinyn®.



Bleed and treatment infusion records

For help with your infusions, please speak to your local hemophilia treatment center or healthcare provider.

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Place label here
and fill out the
adjacent details

Place label here
and fill out the
adjacent details

Date: _____

Location of bleed: _____

Cause of bleed: _____

Duration of bleed: _____

Details: _____

Please see additional Important Safety Information on page 11. Please see Prescribing Information [here](#).

This logbook may last for a longer period of time, based upon your number of bleeds.

rebinyn®

Coagulation Factor IX
(Recombinant), GlycoPEGylated

Important Safety Information

Indications and Usage

What is Rebinyn®, Coagulation Factor IX (Recombinant), GlycoPEGylated?

Rebinyn® is an injectable medicine used to replace clotting Factor IX that is missing in patients with hemophilia B. Rebinyn® is used to treat, prevent, or reduce the frequency (number) of bleeding episodes in people with hemophilia B. Your healthcare provider may give you Rebinyn® when you have surgery. Rebinyn® is not used for immune tolerance therapy.

Important Safety Information

What is the most important information I need to know about Rebinyn®?

- **Do not attempt to do an infusion yourself unless you have been taught how by your healthcare provider or hemophilia treatment center.** Carefully follow your healthcare provider's instructions regarding the dose and schedule for infusing Rebinyn®.

Who should not use Rebinyn®?

Do not use Rebinyn® if you:

- are allergic to Factor IX or any of the other ingredients of Rebinyn®.
- are allergic to hamster proteins.

What should I tell my healthcare provider before using Rebinyn®?

Tell your healthcare provider if you:

- have or have had any medical conditions.
- take any medicines, including non-prescription medicines and dietary supplements.
- are nursing, pregnant, or plan to become pregnant.
- have been told you have inhibitors to Factor IX.

How should I use Rebinyn®?

- Rebinyn® is given as an infusion into the vein.
- **Call your healthcare provider right away if your bleeding does not stop after taking Rebinyn®.**
- Do not stop using Rebinyn® without consulting your healthcare provider.

What are the possible side effects of Rebinyn®?

- **Common side effects include** infusion site reaction (bruising, bleeding, swelling, pain, or redness), itching, and rash.
- Your body can also make antibodies called "inhibitors" against Factor IX, including Rebinyn®, which may stop Rebinyn® from working properly. Your healthcare provider may need to test your blood for inhibitors from time to time.
- **Call your healthcare provider right away or get emergency treatment right away if you get, for example, any of the following signs of an allergic reaction:** hives, chest tightness, wheezing, difficulty breathing, and/or swelling of the face.
- You may be at an increased risk of forming blood clots in your body, especially if you have risk factors for developing blood clots. Call your healthcare provider if you have chest pain, difficulty breathing, leg tenderness, or swelling.
- Animals given repeat doses of Rebinyn® showed Polyethylene Glycol (PEG) in certain cells in the brain. The potential human implications of these animal tests are unknown.

Rebinyn® is a prescription medication.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

BE READY FOR THE UNEXPECTED



Timothy has severe hemophilia B and uses Rebinyn®.

► Get the answers you need

Being proactive in your treatment process requires reaching out for information and support when you need it. The following actions can help you take control of your Rebinyn® treatment when you have questions.

- 1 | Speak to your hemophilia treatment center or healthcare provider for any questions regarding infusions.
- 2 | Visit **Rebinyn.com** to stay informed on recent updates and the latest product information.
- 3 | Contact your local Novo Nordisk Hemophilia Community Liaison for additional information.

Please see additional Important Safety Information on page 11.
Please see Prescribing Information [here](#).



Novo Nordisk Inc., 800 Scudders Mill Road, Plainsboro, New Jersey 08536 U.S.A.

MixPro® and Rebinyn® are registered trademarks and NovoSecure™ is a trademark of Novo Nordisk Health Care AG.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2022 Novo Nordisk Printed in the U.S.A. US22REB00171 October 2022

rebinyn®
*Coagulation Factor IX
(Recombinant), GlycoPEGylated*